

Extended theory of harmonic maps connects general relativity to chaos and quantum mechanism

Gang Ren ^{a,*}, Yi-Shi Duan ^{b,*}

^a The Molecular Foundry, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley CA 94720, USA

^b The Department of Physics, Lanzhou University, Lanzhou, Gansu 730000, China

ARTICLE INFO

Article history:

Received 16 September 2016

Revised 11 March 2017

Accepted 6 July 2017

Keywords:

Extended harmonic maps
Extended theory of harmonic maps
Chaos equation
General relativity
Quantum physics

ABSTRACT

General relativity and quantum mechanism are two separate rules of modern physics explaining how nature works. Both theories are accurate, but the direct connection between two theories was not yet clarified. Recently, researchers blur the line between classical and quantum physics by connecting chaos and entanglement equation. Here, we showed the Duan's extended HM theory, which has the solution of the general relativity, can also have the solutions of the classic chaos equations and even the solution of Schrödinger equation in quantum physics, suggesting the extended theory of harmonic maps may act as a universal theory of physics.

Published by Elsevier Ltd.

This is an open access article under the CC BY license. (<http://creativecommons.org/licenses/by/4.0/>)

1. Introduction

Harmonic map (HM) has been used to recover the Ernst formulation of Einstein's equations in general relativity [1–4], in which the solution of Euler-Lagrange equations of the harmonic maps could be obtained through solving the Laplace-Beltrami's equation and the geodesic equations [3]. From the viewpoint of physics, the traditional theory of harmonic maps contains only the kinetic energy term. For a wider application of this method, the Lagrangian of the traditional HM theory was supplemented with a potential energy term by Duan [5]. This extended HM theory is helpful for studying the travelling wave solutions of some types of nonlinear partial differential equations. Recently, researchers discovered the classical chaos showed a connection to quantum physics via entanglement [6,7]. Here, we asked whether the extended HM theory can recover the classical chaos questions even the Schrödinger equation in quantum physics.

In classic physics, all physical processes are described by differential equations. Over the last several decades, many nonlinear ordinary differential equations possessing chaotic behaviours have been discovered [8], including the nonlinear equation for anharmonic system in periodic fields [9] given by

$$\frac{d^2x}{dt^2} + k \frac{dx}{dt} - \beta x + \alpha x^3 = b \cos(\varpi t), \quad (1)$$

where k , α and β are control parameters, and equation for a parametrically excited pendulum given by [10]

$$\frac{d^2x}{dt^2} + k \frac{dx}{dt} + [\alpha + \beta \cos(\varpi t)] \sin(x) = 0, \quad (2)$$

where k , α and β are control parameters. To date, the study of chaotic behaviours using the space distribution of nonlinear partial differential equations has been rare and is currently only in its initial stage.

Here, we used the extended theory of HM, as the express of the space distribution of nonlinear partial differential equations, derivate the classical chaos Eqs. (1) and (2). Considering the chaotic behaviours connects to quantum mechanism via quantum engagement [6,7], we further found the extended theory of HM has the solution of the Schrödinger equation for a one-dimensional harmonic oscillator. Considering the extended theory of HM has solutions of the Einstein's equations, classic chaotic solutions and Schrödinger equation, suggesting the extended theory of HM may function as a fundamental rule our universe of physics.

2. The extended theory of harmonic maps (HM)

The theory of HM [11–13] became an important branch of mathematical physics decades ago and has been applied to a wide variety of problems in mathematics and theoretical physics. In this section, we re-introduce the formulation of the extended theory of HM that has been reported by Duan [5].

Let M and N be two Riemannian manifolds with local coordinates x^μ ($\mu = 1, 2, \dots, m$) on M and local coordinates

* Corresponding authors.

E-mail addresses: gren@lbl.gov, garyren@yahoo.com (G. Ren), ysduan@lzu.edu.cn (Y.-S. Duan).

Φ^A ($A = 1, 2, \dots, n$) on N . The metrics on M and N are denoted by

$$\begin{aligned} dI^2 &= g_{\mu\nu}(x)dx^\mu dy^\nu; \quad \dim(M) = m \\ dL^2 &= G_{AB}(\Phi)d\Phi^A d\Phi^B; \quad \dim(N) = n \end{aligned} \quad (3)$$

respectively. A mapping

$$\begin{aligned} \Phi: M &\rightarrow N \\ x &\rightarrow \Phi(x) \end{aligned}$$

is called an extended harmonic map if it satisfies the Euler–Lagrange equation resulting from the variational principle $\delta I = 0$, using the action

$$I = \int d^nx \sqrt{g} \left[-\frac{1}{2} g^{\mu\nu} \partial_\mu \Phi^A \partial_\nu \Phi^B G_{AB}(\Phi) + V(\Phi) \right] \quad (4)$$

where $\partial_\mu = \frac{\partial}{\partial x^\mu}$, $V(\Phi)$ is the potential function of Φ^A , and $V(\Phi) = V(\Phi^1, \Phi^2, \dots, \Phi^n)$.

It is obvious that the traditional HM corresponds to the case of $V(\Phi) = 0$ in Eq. (4). It has been showed that, in two dimensional case, the solutions of the Ernst equation in Einstein's general relativity can be recovered from the Euler–Lagrange's equations of traditional HM [3,14].

The conditions for a map to be harmonic are given by the Euler–Lagrange equations

$$\frac{\partial L}{\partial \Phi^A} - \partial_\mu \frac{\partial L}{\partial \partial_\mu x^\Phi} = 0; \quad A = 1, 2, \dots, n \quad (5)$$

where

$$L = -\frac{1}{2} g^{\mu\nu} \partial_\mu \Phi^A \partial_\nu \Phi^B G_{AB}(\Phi) \sqrt{g} + V(\Phi) \sqrt{g} \quad (6)$$

The Eq. (5) follows the notational conventions of Wald's equation [15]. This Eq. (5) appeared in the original paper of Duan (as Eq. 2) [5].

By substituting (6) into (5), we can obtain the Euler–Lagrange equations of extended HM given by

$$\frac{1}{\sqrt{g}} \partial_\mu (\sqrt{g} g^{\mu\nu} \partial_\nu \Phi^A) + \Gamma_{BC}^A \partial_\mu \Phi^B \partial_\nu \Phi^C g^{\mu\nu} + G^{AB} \frac{\partial V(\Phi)}{\partial \Phi^B} = 0, \quad (7)$$

where Γ_{BC}^A are the Christoffels symbols on manifold N given by

$$\Gamma_{BC}^A = \frac{1}{2} G^{AD} \left[\frac{\partial G_{BD}}{\partial \Phi^C} + \frac{\partial G_{CD}}{\partial \Phi^B} - \frac{\partial G_{BC}}{\partial \Phi^D} \right]. \quad (8)$$

In order to obtain a special type of solution of partial differential Eq. (7), Duan used his published approach [3], which could be especial convenient to be used to study the solution of the partial differential Eq. (7). In brief, in the case of Φ^A ($A = 1, 2, \dots, n$) are functions solely of the argument σ , and σ is a function of x^μ on the manifold M :

$$\Phi^A = \Phi^A(\sigma); \quad (9)$$

$$\sigma = \sigma(x). \quad (10)$$

The Eq. (7) can be written as following,

$$\begin{aligned} \frac{1}{\sqrt{g}} \partial_\mu (\sqrt{g} g^{\mu\nu} \partial_\nu \sigma) \frac{\partial \Phi^A}{\partial \sigma} + G^{AB} \frac{\partial V(\Phi)}{\partial \Phi^B} \\ + \left(\frac{d^2 \Phi^A}{d\sigma^2} + \Gamma_{BC}^A \frac{d\Phi^B}{d\sigma} \frac{d\Phi^C}{d\sigma} \right) g^{\mu\nu} \partial_\mu \sigma \partial_\nu \sigma = 0. \end{aligned} \quad (11)$$

This important Eq. (11) appear in the original paper of Duan (as Eq. 7) [5].

The solution, such as the solution of Sine-Gorden equation, and the travelling wave solution can be found in certain cases of the metrics and potential function in the extended Euler–Lagrange Eq. (11).

If the function $\sigma = \sigma(x)$ satisfies the Laplace–Beltrami equations

$$\frac{1}{\sqrt{g}} \partial_\mu (\sqrt{g} g^{\mu\nu} \partial_\nu \sigma) = 0, \quad (12)$$

the Eq. (11) will take the form

$$\frac{d^2 \Phi^A}{d\sigma^2} + \Gamma_{BC}^A \frac{d\Phi^B}{d\sigma} \frac{d\Phi^C}{d\sigma} = -\frac{1}{f} G^{AB} \frac{\partial V(\Phi)}{\partial \Phi^B}, \quad (13)$$

where

$$f = g^{\mu\nu} \partial_\mu \sigma \partial_\nu \sigma \quad (14)$$

is a scalar function on the manifold M .

Because f is not the function of Φ^A , a rescaling of $V(\Phi)$ as $V(\Phi) = fU(\Phi)$ re-expresses the Eq. (13) as

$$\frac{d^2 \Phi^A}{d\sigma^2} + \Gamma_{BC}^A \frac{d\Phi^B}{d\sigma} \frac{d\Phi^C}{d\sigma} = -G^{AB} \frac{\partial U(\Phi)}{\partial \Phi^B}. \quad (15)$$

Eq. (15) can be represented as the geodesic equation of a particle on the Riemannian manifold N subjected to an external force of

$$F^A = -G^{AB} \frac{\partial U(\Phi)}{\partial \Phi^B}. \quad (16)$$

This Eq. (15) will become equivalent to the usual geodesic equations on the manifold N if the external force $F^A = 0$ ($A = 1, 2, \dots, n$). Eq. (15) is physical dynamical equation.

3. Chaotic solutions of the extended HM equations

If the chaos equation could be derive from the Eq. (15) under certain cases of metrics and potential function, the chaotic behaviour satisfied the extended HM equations. In order to evidence whether the chaotic behaviours exist in the partial differential Eq. (15), we simplified the case as that the M is the pseudo-Euclidean space-time and N is a 2-dimensional manifold with coordinates $\{\Phi^1, \Phi^2\}$. We suppose that Φ^1 and Φ^2 are functions of an argument $\sigma = \sigma(x) = k_\mu x^\mu$ and further assume that $\Phi^2 = \sigma$, i.e.,

$$\Phi^1 = \Phi(\sigma); \quad \Phi^2 = \sigma; \quad \sigma = k_\mu x^\mu. \quad (17)$$

Under this simplified case, the Eq. (15) can be expressed as

$$\begin{aligned} \frac{d^2 \Phi}{d\sigma^2} + \Gamma_{11}^1 \left(\frac{d\Phi}{d\sigma} \right)^2 + 2\Gamma_{12}^1 \frac{d\Phi}{d\sigma} + \Gamma_{22}^1 \\ = - \left(G^{11} \frac{\partial U(\Phi, \sigma)}{\partial \Phi} + G^{12} \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right); \end{aligned} \quad (18)$$

$$\begin{aligned} \Gamma_{11}^2 \left(\frac{d\Phi}{d\sigma} \right)^2 + 2\Gamma_{12}^2 \frac{d\Phi}{d\sigma} + \Gamma_{22}^2 \\ = - \left(G^{21} \frac{\partial U(\Phi, \sigma)}{\partial \Phi} + G^{22} \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right). \end{aligned} \quad (19)$$

Eliminating the term $(\frac{d\Phi}{d\sigma})^2$ from the differential Eqs. (18) and (19), we obtain

$$\begin{aligned} \frac{d^2 \Phi}{d\sigma^2} + 2 \left(\Gamma_{12}^1 - \frac{\Gamma_{11}^1 \Gamma_{12}^2}{\Gamma_{11}^2} \right) \frac{d\Phi}{d\sigma} + \left(\Gamma_{22}^1 - \frac{\Gamma_{11}^1 \Gamma_{22}^2}{\Gamma_{11}^2} \right) \\ + \left[G^{11} \frac{\partial U(\Phi, \sigma)}{\partial \Phi} + G^{12} \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right. \\ \left. - \frac{\Gamma_{11}^1}{\Gamma_{11}^2} \left(G^{21} \frac{\partial U(\Phi, \sigma)}{\partial \Phi} + G^{22} \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right) \right] = 0 \end{aligned} \quad (20)$$

If the chaos equation can be derive from these geodesical ordinary differential equations (20), the partial differential equation (15) must contain the chaotic behaviours.

By given the metrics on manifold N are diagonal and take the following form:

$$G_{11} = e^{\Phi+\sigma}; \quad G_{12} = 0; \quad G_{21} = 0; \quad G_{22} = (k-1)e^{\Phi+\sigma}, \quad (21)$$

where k is a constant.

The Christoffels symbols (8) on the 2-dimensional manifold are calculated as following,

$$\begin{aligned} \Gamma_{11}^1 &= \frac{1}{2}; \quad \Gamma_{12}^1 = \frac{1}{2}; \quad \Gamma_{22}^1 = -\frac{1}{2}(k-1); \\ \Gamma_{11}^2 &= -\frac{1}{2(k-1)}; \quad \Gamma_{12}^2 = \frac{1}{2}; \quad \Gamma_{22}^2 = \frac{1}{2}. \end{aligned} \quad (22)$$

The detailed procedures of calculation showed in the supplementary information.

By substituting (21) and (22) into (20), the differential Eq. (20) can be written as following,

$$\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + (e^{-\Phi-\sigma}) \left[\frac{\partial U(\Phi, \sigma)}{\partial \Phi} + \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] = 0. \quad (23)$$

We found, under following given potential function $U(\Phi, \sigma)$, i.e.

$$\begin{aligned} U(\Phi, \sigma) &= e^{\Phi+\sigma} \left[\frac{1}{2}\alpha\Phi^3 - \frac{3}{4}\alpha\Phi^2 - \frac{1}{4}(2\beta - 3\alpha)\Phi \right. \\ &\quad + \frac{1}{8}(2\beta - 3\alpha) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) \\ &\quad \left. - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right], \end{aligned} \quad (24)$$

Substituting (24) into Eq. (23), we obtain

$$\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} - \beta\Phi + \alpha\Phi^3 = b\cos(\omega\sigma); \quad \sigma = k_\mu x^\mu \quad (25)$$

The detailed procedures of calculation showed in the supplementary information.

Comparing the Eq. (25) with (1), i.e., the equation of anharmonic system in a periodic field, we find that they are of the same form. Since the Eq. (1) possesses chaotic states that are characterized by the existence of a strange attractor in the phase space, this implies that Eq. (23) for $\Phi = \Phi(\sigma)$ should possess the same chaotic profile.

By giving a new potential function $U = U(\Phi, \sigma)$, i.e.

$$\begin{aligned} U(\Phi, \sigma) &= e^{\Phi+\sigma} \left\{ \frac{2}{5}\alpha \sin(\Phi) - \frac{1}{5}\alpha \cos(\Phi) \right. \\ &\quad + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) \\ &\quad - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) \\ &\quad \left. + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi) \right\}, \end{aligned} \quad (26)$$

in which, α and β are constants, the Eq. (23) become following,

$$\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + [\alpha + \beta \cos(\omega\sigma)] \sin(\Phi) = 0; \quad \sigma = k_\mu x^\mu \quad (27)$$

The detailed procedures of calculation showed in the supplementary information.

Comparing the Eq. (27) with (2), i.e., the nonlinear equation of a parametrically excited damped pendulum, we find that they are of the same form. Since the Eq. (2) possesses chaotic states that are characterized by the existence of a strange attractor in the phase space, this implies that Eq. (27) for $\Phi(\sigma)$ should possess the same chaotic profile.

4. Quantum mechanism solutions of the extended HM equations

The extended equation of HM contains the chaotic solution, which recently found with connection to quantum physics via entanglement [1,2]. It would be interested to test whether the Schrodinger equation in quantum physics can be derived. As a simple case, we studied the Schrodinger equation for a one-dimensional harmonic oscillator as below,

$$\frac{d^2\Psi}{dx^2} = -\frac{2m}{\hbar^2} \left(E - \frac{1}{2}Kx^2 \right) \Psi(x), \quad (28)$$

where $\Psi(x)$ is the wave function, E is the total energy (constant), and K is the force constant (the force on the mass being $F = -Kx$, proportional to the displacement x and directed towards the origin).

By given the metrics on manifold N are diagonal and take the following form:

$$G_{11} = e^{\Phi+\sigma}; \quad G_{12} = 0; \quad G_{21} = 0; \quad G_{22} = -e^{\Phi+\sigma}, \quad (29)$$

This Eq. (29) is a special case of Eq. (21), i.e. $k = 0$. Thus, the differential Eq. (20) can be written as following,

$$\frac{d^2\Phi}{d\sigma^2} + (e^{-\Phi-\sigma}) \left[\frac{\partial U(\Phi, \sigma)}{\partial \Phi} + \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] = 0. \quad (30)$$

By given the potential function $U(\Phi, \sigma)$ as following, i.e.

$$\begin{aligned} U(\Phi, \sigma) &= e^{\Phi+\sigma} \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 - \frac{m}{4\hbar^2} K\sigma^2 - \frac{m}{2\hbar^2} K\Phi\sigma \right. \\ &\quad \left. + \frac{2m}{\hbar^2} (E - K)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{\hbar^2} \left(\frac{3}{4}K - E \right) \right], \end{aligned} \quad (31)$$

where E , K , m , \hbar are the constants,

Substituting (31) into Eq. (30), we obtain

$$\frac{d^2\Phi}{d\sigma^2} = -\frac{2m}{\hbar^2} \left(E - \frac{1}{2}K\sigma^2 \right) \Phi(\sigma), \quad (32)$$

The detailed procedures of calculation showed in the supplementary information.

Comparing the Eq. (32) with Eq. (28), we find that they are of the same form when $\Phi(\sigma) = \Psi(x)$ and $\sigma = x$. Since the Eq. (28) is one of typical equation of Schrodinger equation in quantum physics, this implies that the Eq. (15) contains the Schrodinger equation in quantum physics.

Since the extended HM (when the potential function $V(\Phi) = 0$ in the Eq. (4)) can be recovered the Einstein's equations in general relativity [14], and the same theory can also recover the Schrodinger equation in quantum physics, we propose here the extended equation of HM as a single and universal theory for our universe.

Author contributions

This project was initiated by GR and YD., GR calculated the solution, and YD. validated the solution. GR drafted the initial manuscript, which was revised by YD.

Competing financial interests

The author(s) declare no competing financial interests.

Acknowledgments


We thank Mr. Wencheng Ren to preserve the original results and manuscript. Work at the Molecular Foundry was supported by the Office of Science, and the Office of Basic Energy Sciences, the U.S. Department of Energy, contract no. DE-AC02-05CH11231.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.chaos.2017.07.004](https://doi.org/10.1016/j.chaos.2017.07.004).

References


- [1] Nguyen L. Singular harmonic maps and applications to general relativity. *Commun Math Phys* 2011;301(2):411–41.
- [2] Beheshti S, Tahvildar-Zadeh S. Integrability and vesture for harmonic maps into symmetric spaces. *Rev Math Phys* 2016;28(3).
- [3] Duan YS. Solution of the Euler's equation of harmonic maps and its application to general relativity. *Acta Phys Sinica* 1984;33.
- [4] Sanchez N. Connection between the nonlinear σ model and the Einstein equations of general relativity. *Phys Rev D* 1982(26).
- [5] Duan YS. Extended harmonic maps equations and the soliton solutions. *J Lanzhou Univ* 1991 S1.
- [6] Neill C, et al. Ergodic dynamics and thermalization in an isolated quantum system. *Nat Phys* 2016.
- [7] Chaudhury S, et al. Quantum signatures of chaos in a kicked top. *Nature* 2009;461(7265):768–71.
- [8] Hao BL. Bifurcation, chaos, strange attractor, turbulence and all that. *Prog Phys* 1983;3.
- [9] Huberman BA, Crutchfield JP. Chaotic states of anharmonic systems in periodic fields. *Phys Rev Lett* 1979;43(23):1743–7.
- [10] Leven RW, Koch BP. Chaotic behavior of a parametrically excited damped pendulum. *Phys Lett A* 1981;86(2):71–4.
- [11] Misner CW. Harmonic maps as models for physical theories. *Phys Rev D* 1978;18.
- [12] Nutku Y, Halil M. Colliding impulsive gravitational-waves. *Phys Rev Lett* 1977;39(22):1379–82.
- [13] Eris A, Nutku Y. Harmonic mappings of Riemannian manifolds and stationary vacuum space-times with whole cylinder symmetry. *J Math Phys* 1975;16(7):1431–4.
- [14] Duan, Y.S., Harmonic maps and their application to general relativity. High-Energy Physics Literature Database, 1983, 194698(slac-pub-3265.pdf).
- [15] Wald RM. General relativity. Chicago: The University of Chicago Press; 1984. p. 455.



Contents lists available at ScienceDirect

Chaos, Solitons and Fractals

Nonlinear Science, and Nonequilibrium and Complex Phenomena

journal homepage: www.elsevier.com/locate/chaos

Corrigendum

Corrigendum to “Extended theory of harmonic maps connects general relativity to chaos and quantum mechanism” [Chaos, Solitons and Fractals 103 (2017) 567–570]

Xiaobo Zhai, Gang Ren*

The Molecular Foundry, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

Abstract

In this corrigendum, we corrected the potential function $U(\Phi, \sigma)$ that used to derive the Schrödinger equation for a one-dimensional harmonic oscillator in [Chaos, Solitons and Fractals 103 (2017) 567–570]. We also corrected several typos in the equations in the paper and the supplementary information. These corrections do not alter the conclusion of the article.

Keywords

Extended harmonic maps; Chaos equation; General relativity; Quantum physics.

1. The definition of the potential function U in Eq. (31)

The definition of the potential function U that used to derive the Schrödinger equation for a one-dimensional harmonic should be

$$U(\Phi, \sigma) = e^{\Phi+\sigma} \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) \right] \quad (1)$$

instead of

$$U(\Phi, \sigma) = e^{\Phi+\sigma} \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 - \frac{m}{4\hbar^2} K\sigma^2 - \frac{m}{2\hbar^2} K\Phi\sigma + \frac{2m}{\hbar^2} (E - K)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{\hbar^2} \left(\frac{3}{4}K - E\right) \right] \quad (2)$$

in Eq. (31) in [1].

Moreover, the Eq. (2) that was also showed in the 2nd equation in the 15th page of supplementary information should be removed.

2. Missed a “K” in the definition of the potential function U of the supplementary information (line #4 in page #15)

The equation in line #4 of page #15 of the supplementary information [1] should be

$$U(\Phi, \sigma) = e^{\Phi+\sigma} \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) \right] \quad (3)$$

instead of

$$U(\Phi, \sigma) = e^{\Phi+\sigma} \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} \sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) \right] \quad (4)$$

DOI of original article: [10.1016/j.chaos.2017.07.004](https://doi.org/10.1016/j.chaos.2017.07.004)

* Corresponding author.

E-mail address: gren@lbl.gov (G. Ren).

3. The typos in Eq. (5)

The Eq. (5) should be

$$\frac{\partial L}{\partial \Phi^A} - \partial_\mu \frac{\partial L}{\partial \partial_\mu \Phi^A} = 0; A = 1, 2, \dots, n \quad (5)$$

instead of

$$\frac{\partial L}{\partial \Phi^A} - \partial_\mu \frac{\partial L}{\partial \partial_\mu x \Phi^A} = 0; A = 1, 2, \dots, n \quad (6)$$

4. The typos in the equation of supplementary information (line #3 in page #2)

In the line #3 of page #2 of supplementary information [1], for the Christoffels symbol Γ_{12}^2 should be

$$\begin{aligned} \Gamma_{12}^2 &= \frac{1}{2} G^{2D} \left[\frac{\partial G_{1D}}{\partial \Phi^2} + \frac{\partial G_{2D}}{\partial \Phi^1} - \frac{\partial G_{12}}{\partial \Phi^D} \right] \\ &= \frac{1}{2} G^{21} \left[\frac{\partial G_{11}}{\partial \sigma} + \frac{\partial G_{21}}{\partial \Phi} - \frac{\partial G_{12}}{\partial \Phi} \right] + \frac{1}{2} G^{22} \left[\frac{\partial G_{12}}{\partial \sigma} + \frac{\partial G_{22}}{\partial \Phi} - \frac{\partial G_{12}}{\partial \sigma} \right] \\ &= \frac{1}{2} G^{22} \left[\frac{\partial G_{22}}{\partial \Phi} \right] = \frac{1}{2} \left(\frac{1}{k-1} e^{-\Phi-\sigma} \right) \left[\frac{\partial[(k-1)e^{\Phi+\sigma}]}{\partial \Phi} \right] = \frac{1}{2} \end{aligned} \quad (7)$$

instead of

$$\begin{aligned} \Gamma_{12}^2 &= \frac{1}{2} G^{2D} \left[\frac{\partial G_{1D}}{\partial \Phi^2} + \frac{\partial G_{2D}}{\partial \Phi^1} - \frac{\partial G_{12}}{\partial \Phi^D} \right] \\ &= \frac{1}{2} G^{21} \left[\frac{\partial G_{11}}{\partial \Phi} + \frac{\partial G_{21}}{\partial \Phi} - \frac{\partial G_{12}}{\partial \Phi} \right] + \frac{1}{2} G^{22} \left[\frac{\partial G_{12}}{\partial \Phi} + \frac{\partial G_{22}}{\partial \Phi} - \frac{\partial G_{12}}{\partial \sigma} \right] \\ &= \frac{1}{2} G^{22} \left[\frac{\partial G_{22}}{\partial \Phi} \right] = \frac{1}{2} \left(\frac{1}{k-1} e^{-\Phi-\sigma} \right) \left[\frac{\partial[(k-1)e^{\Phi+\sigma}]}{\partial \Phi} \right] = \frac{1}{2} \end{aligned} \quad (8)$$

5. The differential symbol of the equation of supplementary information (line #10 in page #6)

The equation in line #10 of page #6 of the supplementary information [1] should be

$$\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \alpha \Phi^3 - \frac{1}{2} (2\beta) \Phi - \frac{4b + b\omega^2}{\omega^2 + 4} \cos(\omega\sigma) = 0 \quad (9)$$

instead of

$$\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \alpha \Phi^3 - \frac{1}{2} (2\beta) \Phi - \frac{4b + b\omega^2}{\omega^2 + 4} \cos(\omega\sigma) = 0 \quad (10)$$

6. The cosine symbol of the equation of supplementary information (line #4 in page #14)

The equation in line #4 of page #14 of the supplementary information [1] should be

$$\begin{aligned} \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25 + 6\omega^2 + \omega^4} \left[-(\omega^2 + 7)\omega \sin(\omega\sigma) \sin(\Phi) - (\omega^2 - 1)\omega \sin(\omega\sigma) \cos(\Phi) + [\omega^4 + 5\omega^2 + 10] \right. \right. \\ \left. \left. \cos(\omega\sigma) \sin(\Phi) - (3\omega^2 + 5)\cos(\omega\sigma) \cos(\Phi) + (\omega^3 + 7\omega)\sin(\omega\sigma) \sin(\Phi) + (\omega^3 - \omega)\sin(\omega\sigma) \cos(\Phi) \right. \right. \\ \left. \left. + (\omega^2 + 15)\cos(\omega\sigma) \sin(\Phi) + (3\omega^2 + 5)\cos(\omega\sigma) \cos(\Phi) + \frac{2}{5}\alpha \sin(\Phi) - \frac{1}{5}\alpha \cos(\Phi) \right] \right] = 0 \end{aligned} \quad (11)$$

instead of

$$\begin{aligned} \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25 + 6\omega^2 + \omega^4} \left[-(\omega^2 + 7)\omega \sin(\omega\sigma) \sin(\Phi) - (\omega^2 - 1)\omega \sin(\omega\sigma) \cos(\Phi) + \right. \right. \\ \left. \left. + [\omega^4 + 5\omega^2 + 10]\cos(\omega\sigma) \sin(\Phi) - (3\omega^2 + 5)\cos(\omega\sigma) \cos(\Phi) + (\omega^3 + 7\omega)\sin(\omega\sigma) \sin(\Phi) + (\omega^3 - \omega)\sin(\omega\sigma) \cos(\Phi) \right. \right. \\ \left. \left. + (\omega^2 + 15)\cos(\omega\sigma) \sin(\Phi) + (3\omega^2 + 5)\cos(\omega\sigma) \cos(\Phi) + \frac{2}{5}\alpha \sin(\Phi) - \frac{1}{5}\alpha \cos(\Phi) \right] \right] = 0 \end{aligned} \quad (12)$$

7. The cosine symbol of the equation of supplementary information (line #11 in page #14)

The equation in line #11 of page #14 of the supplementary information [1] should be

$$\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25 + 6\omega^2 + \omega^4} \left[-(\omega^2 + 7)\omega \sin(\omega\sigma) \sin(\Phi) + (\omega^3 + 7\omega) \sin(\omega\sigma) \sin(\Phi) \right. \right. \\ \left. \left. - (\omega^2 - 1)\omega \sin(\omega\sigma) \cos(\Phi) + (\omega^3 - \omega) \sin(\omega\sigma) \cos(\Phi) + [\omega^4 + 5\omega^2 + 10] \cos(\omega\sigma) \sin(\Phi) \right. \right. \\ \left. \left. + (\omega^2 + 15) \cos(\omega\sigma) \sin(\Phi) - (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi) + (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi) \right] \right] = 0 \quad (13)$$

instead of

$$\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25 + 6\omega^2 + \omega^4} \left[-(\omega^2 + 7)\omega \sin(\omega\sigma) \sin(\Phi) + (\omega^3 + 7\omega) \sin(\omega\sigma) \sin(\Phi) \right. \right. \\ \left. \left. - (\omega^2 - 1)\omega \sin(\omega\sigma) \cos(\Phi) + (\omega^3 - \omega) \sin(\omega\sigma) \cos(\Phi) + [\omega^4 + 5\omega^2 + 10] \cos(\omega\sigma) \sin(\Phi) \right. \right. \\ \left. \left. + (\omega^2 + 15) \cos(\omega\sigma) \sin(\Phi) - (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi) + (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi) \right] \right] = 0 \quad (14)$$

8. The cosine symbol of the equation of supplementary information (line #13 in page #14)

The equation in line #13 of page #14 of the supplementary information [1] should be

$$\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25 + 6\omega^2 + \omega^4} [(\omega^4 + 5\omega^2 + 10 + \omega^2 + 15) \cos(\omega\sigma) \sin(\Phi)] \right] = 0 \quad (15)$$

instead of

$$\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25 + 6\omega^2 + \omega^4} [(\omega^4 + 5\omega^2 + 10 + \omega^2 + 15) \cos(\omega\sigma) \sin(\Phi)] \right] = 0 \quad (16)$$

9. The symbol of the equation of supplementary information (line #4 in page #17)

The equation in line #4 of page #17 of the supplementary information [1] should be

$$\frac{\partial U(\Phi, \sigma)}{\partial \sigma} = e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) \right. \\ \left. + \frac{\partial \left[-\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) \right]}{\partial \sigma} \right\} \\ = e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) + \frac{\partial \left[-\frac{m}{2\hbar^2} K \Phi \sigma^2 \right]}{\partial \sigma} \right. \\ \left. + \frac{\partial \left[\frac{m}{4\hbar^2} K \sigma^2 \right]}{\partial \sigma} + \frac{\partial \left[\frac{m}{2\hbar^2} K \Phi \sigma \right]}{\partial \sigma} + \frac{\partial \left[\frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right) \Phi \right]}{\partial \sigma} + \frac{\partial \left[-\frac{m}{2\hbar^2} K \sigma \right]}{\partial \sigma} + \frac{\partial \left[\frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) \right]}{\partial \sigma} \right\} \\ = e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) - \frac{m}{2\hbar^2} K \Phi \frac{\partial [\sigma^2]}{\partial \sigma} \right. \\ \left. + \frac{m}{4\hbar^2} K \frac{\partial [\sigma^2]}{\partial \sigma} + \frac{m}{2\hbar^2} K \Phi \frac{\partial [\sigma]}{\partial \sigma} + 0 - \frac{m}{2\hbar^2} K \frac{\partial [\sigma]}{\partial \sigma} + 0 \right\} \\ = e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) - \frac{m}{\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} K \sigma \right. \\ \left. + \frac{m}{2\hbar^2} K \Phi - \frac{m}{2\hbar^2} K \right\} \\ = e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma - \frac{m}{\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right) \Phi + \frac{m}{2\hbar^2} K \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} K \sigma \right. \\ \left. + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) - \frac{m}{2\hbar^2} K \right\} \\ = e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 - \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} \left(2E + \frac{1}{2}K\right) \Phi - \frac{m}{2\hbar^2} \left(\frac{1}{4}K + E\right) \right\} \quad (17)$$

instead of

$$\begin{aligned}
 \frac{\partial U(\Phi, \sigma)}{\partial \sigma} &= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) \right. \\
 &\quad \left. + \frac{\partial \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) \right]}{\partial \sigma} \right\} \\
 &= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) + \frac{\partial \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 \right]}{\partial \sigma} \right. \\
 &\quad \left. + \frac{\partial \left[\frac{m}{4\hbar^2} K\sigma^2 \right]}{\partial \sigma} + \frac{\partial \left[\frac{m}{2\hbar^2} K\Phi\sigma \right]}{\partial \sigma} + \frac{\partial \left[\frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi \right]}{\partial \sigma} + \frac{\partial \left[-\frac{m}{2\hbar^2} K\sigma \right]}{\partial \sigma} + \frac{\partial \left[\frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) \right]}{\partial \sigma} \right\} \\
 &= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) - \frac{m}{2\hbar^2} K\Phi \frac{\partial [\sigma^2]}{\partial \sigma} \right. \\
 &\quad \left. + \frac{m}{4\hbar^2} K \frac{\partial [\sigma^2]}{\partial \sigma} + \frac{m}{2\hbar^2} K\Phi \frac{\partial [\sigma]}{\partial \sigma} + 0 - \frac{m}{2\hbar^2} K \frac{\partial [\sigma]}{\partial \sigma} + 0 \right\} \\
 &= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) - \frac{m}{\hbar^2} K\Phi\sigma \right. \\
 &\quad \left. + \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} K\Phi - \frac{m}{2\hbar^2} K \right\} \\
 &= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma - \frac{m}{\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi + \frac{m}{2\hbar^2} K\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} K\sigma \right. \\
 &\quad \left. + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) - \frac{m}{2\hbar^2} K \right\} \\
 &= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 - \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E + \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} \left(\frac{1}{4}K + E\right) \right\} \tag{18}
 \end{aligned}$$

Reference

[1] Ren G, Duan Y-S. Extended theory of harmonic maps connects general relativity to chaos and quantum mechanism. *Chaos Solitons Fractals* 2017;103:567–70.

For two dimensional: $\Phi^1 = \Phi(\sigma)$; $\Phi^2 = \sigma$, the equation (20) is as below

$$\frac{d^2\Phi}{d\sigma^2} + 2 \left(\Gamma_{12}^1 - \frac{\Gamma_{11}^1 \Gamma_{12}^2}{\Gamma_{11}^2} \right) \frac{d\Phi}{d\sigma} + \left(\Gamma_{22}^1 - \frac{\Gamma_{11}^1 \Gamma_{22}^2}{\Gamma_{11}^2} \right) + \left\{ G^{11} \frac{\partial U(\Phi, \sigma)}{\partial \Phi} + G^{12} \frac{\partial U(\Phi, \sigma)}{\partial \sigma} - \frac{\Gamma_{11}^1}{\Gamma_{11}^2} \left[G^{21} \frac{\partial U(\Phi, \sigma)}{\partial \sigma} + G^{22} \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] \right\} = 0$$

Metric:

$$G_{11} = e^{\Phi+\sigma}; \quad G_{12} = 0; \quad G_{21} = 0; \quad G_{22} = (k-1)e^{\Phi+\sigma}$$

$$G^{11} = e^{-\Phi-\sigma}; \quad G^{12} = 0; \quad G^{21} = 0; \quad G^{22} = \frac{1}{k-1} e^{-\Phi-\sigma}$$

Christoffels symbols:

$$\Gamma_{BC}^A = \frac{1}{2} G^{AD} \left[\frac{\partial G_{BD}}{\partial \Phi^C} + \frac{\partial G_{CD}}{\partial \Phi^B} - \frac{\partial G_{BC}}{\partial \Phi^D} \right]$$

$$\begin{aligned} \Gamma_{11}^1 &= \frac{1}{2} G^{1D} \left[\frac{\partial G_{1D}}{\partial \Phi^1} + \frac{\partial G_{1D}}{\partial \Phi^1} - \frac{\partial G_{11}}{\partial \Phi^D} \right] = \frac{1}{2} G^{11} \left[\frac{\partial G_{11}}{\partial \Phi} + \frac{\partial G_{11}}{\partial \Phi} - \frac{\partial G_{11}}{\partial \sigma} \right] + \frac{1}{2} G^{12} \left[\frac{\partial G_{12}}{\partial \Phi} + \frac{\partial G_{12}}{\partial \Phi} - \frac{\partial G_{11}}{\partial \sigma} \right] = \frac{1}{2} G^{11} \left[\frac{\partial G_{11}}{\partial \Phi} \right] = \frac{1}{2} (e^{\Phi+\sigma}) \left[\frac{\partial (e^{\Phi+\sigma})}{\partial \Phi} \right] \\ &= \frac{1}{2} \\ \Gamma_{12}^1 &= \frac{1}{2} G^{1D} \left[\frac{\partial G_{1D}}{\partial \Phi^2} + \frac{\partial G_{2D}}{\partial \Phi^1} - \frac{\partial G_{12}}{\partial \Phi^D} \right] = \frac{1}{2} G^{11} \left[\frac{\partial G_{11}}{\partial \sigma} + \frac{\partial G_{21}}{\partial \Phi} - \frac{\partial G_{12}}{\partial \Phi} \right] + \frac{1}{2} G^{12} \left[\frac{\partial G_{12}}{\partial \sigma} + \frac{\partial G_{22}}{\partial \Phi} - \frac{\partial G_{12}}{\partial \sigma} \right] = \frac{1}{2} G^{11} \left[\frac{\partial G_{11}}{\partial \sigma} \right] = \frac{1}{2} (e^{-\Phi-\sigma}) \left[\frac{\partial (e^{\Phi+\sigma})}{\partial \sigma} \right] \\ &= \frac{1}{2} \\ \Gamma_{22}^1 &= \frac{1}{2} G^{1D} \left[\frac{\partial G_{2D}}{\partial \Phi^2} + \frac{\partial G_{2D}}{\partial \Phi^2} - \frac{\partial G_{22}}{\partial \Phi^D} \right] = \frac{1}{2} G^{11} \left[\frac{\partial G_{21}}{\partial \sigma} + \frac{\partial G_{21}}{\partial \sigma} - \frac{\partial G_{22}}{\partial \Phi} \right] + \frac{1}{2} G^{12} \left[\frac{\partial G_{22}}{\partial \sigma} + \frac{\partial G_{22}}{\partial \sigma} - \frac{\partial G_{22}}{\partial \sigma} \right] = \frac{1}{2} G^{11} \left[-\frac{\partial G_{22}}{\partial \Phi} \right] \\ &= \frac{1}{2} (e^{-\Phi-\sigma}) \left[-\frac{\partial [(k-1)e^{\Phi+\sigma}]}{\partial \Phi} \right] = -\frac{1}{2} (k-1) \end{aligned}$$

$$\Gamma_{11}^2 = \frac{1}{2} G^{2D} \left[\frac{\partial G_{1D}}{\partial \Phi^1} + \frac{\partial G_{1D}}{\partial \Phi^1} - \frac{\partial G_{11}}{\partial \Phi^D} \right] = \frac{1}{2} G^{21} \left[\frac{\partial G_{11}}{\partial \Phi} + \frac{\partial G_{11}}{\partial \Phi} - \frac{\partial G_{11}}{\partial \sigma} \right] + \frac{1}{2} G^{22} \left[\frac{\partial G_{12}}{\partial \Phi} + \frac{\partial G_{12}}{\partial \Phi} - \frac{\partial G_{11}}{\partial \sigma} \right] = \frac{1}{2} G^{22} \left[-\frac{\partial G_{11}}{\partial \sigma} \right]$$

$$= \frac{1}{2} \left(\frac{1}{k-1} e^{-\Phi-\sigma} \right) \left[-\frac{\partial (e^{\Phi+\sigma})}{\partial \sigma} \right] = -\frac{1}{2(k-1)}$$

$$\Gamma_{12}^2 = \frac{1}{2} G^{2D} \left[\frac{\partial G_{1D}}{\partial \Phi^2} + \frac{\partial G_{2D}}{\partial \Phi^1} - \frac{\partial G_{12}}{\partial \Phi^D} \right] = \frac{1}{2} G^{21} \left[\frac{\partial G_{11}}{\partial \Phi} + \frac{\partial G_{21}}{\partial \Phi} - \frac{\partial G_{12}}{\partial \Phi} \right] + \frac{1}{2} G^{22} \left[\frac{\partial G_{12}}{\partial \Phi} + \frac{\partial G_{22}}{\partial \Phi} - \frac{\partial G_{12}}{\partial \sigma} \right] = \frac{1}{2} G^{22} \left[\frac{\partial G_{22}}{\partial \Phi} \right]$$

$$= \frac{1}{2} \left(\frac{1}{k-1} e^{-\Phi-\sigma} \right) \left[\frac{\partial [(k-1)e^{\Phi+\sigma}]}{\partial \Phi} \right] = \frac{1}{2}$$

$$\Gamma_{22}^2 = \frac{1}{2} G^{2D} \left[\frac{\partial G_{2D}}{\partial \Phi^2} + \frac{\partial G_{2D}}{\partial \Phi^2} - \frac{\partial G_{22}}{\partial \Phi^D} \right] = \frac{1}{2} G^{21} \left[\frac{\partial G_{21}}{\partial \sigma} + \frac{\partial G_{21}}{\partial \sigma} - \frac{\partial G_{22}}{\partial \Phi} \right] + \frac{1}{2} G^{22} \left[\frac{\partial G_{22}}{\partial \sigma} + \frac{\partial G_{22}}{\partial \sigma} - \frac{\partial G_{22}}{\partial \sigma} \right] = \frac{1}{2} G^{22} \left[\frac{\partial G_{22}}{\partial \sigma} \right]$$

$$= \frac{1}{2} \left(\frac{1}{k-1} e^{-\Phi-\sigma} \right) \left[\frac{\partial [(k-1)e^{\Phi+\sigma}]}{\partial \sigma} \right] = \frac{1}{2}$$

The equation (20) can be calculated as

$$\frac{d^2 \Phi}{d\sigma^2} + 2 \left(\Gamma_{12}^1 - \frac{\Gamma_{11}^1 \Gamma_{12}^2}{\Gamma_{11}^2} \right) \frac{d\Phi}{d\sigma} + \left(\Gamma_{22}^1 - \frac{\Gamma_{11}^1 \Gamma_{22}^2}{\Gamma_{11}^2} \right) + \left\{ G^{11} \frac{\partial U(\Phi, \sigma)}{\partial \Phi} + G^{12} \frac{\partial U(\Phi, \sigma)}{\partial \sigma} - \left(\frac{\Gamma_{11}^1}{\Gamma_{11}^2} \right) \left[G^{21} \frac{\partial U(\Phi, \sigma)}{\partial \Phi} + G^{22} \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] \right\} = 0$$

$$\frac{d^2 \Phi}{d\sigma^2} + 2 \left(\Gamma_{12}^1 - \frac{\Gamma_{11}^1 \Gamma_{12}^2}{\Gamma_{11}^2} \right) \frac{d\Phi}{d\sigma} + \left(\Gamma_{22}^1 - \frac{\Gamma_{11}^1 \Gamma_{22}^2}{\Gamma_{11}^2} \right) + \left[G^{11} \frac{\partial U(\Phi, \sigma)}{\partial \Phi} - \left(\frac{\Gamma_{11}^1}{\Gamma_{11}^2} \right) G^{22} \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] = 0$$

in which,

$$\left(\Gamma_{12}^1 - \frac{\Gamma_{11}^1 \Gamma_{12}^2}{\Gamma_{11}^2} \right) = \frac{1}{2} - \frac{\frac{1}{2} * \frac{1}{2}}{-2(k-1)} = \frac{1}{2} + \frac{(k-1)}{2} = \frac{k}{2}$$

$$\left(\Gamma_{22}^1 - \frac{\Gamma_{11}^1 \Gamma_{22}^2}{\Gamma_{11}^2} \right) = -\frac{1}{2} (k-1) - \frac{\frac{1}{2} * \frac{1}{2}}{-2(k-1)} = -\frac{(k-1)}{2} + \frac{(k-1)}{2} = 0$$

$$\left(\frac{\Gamma_{11}^1}{\Gamma_{11}^2}\right) = -\frac{\frac{1}{2}}{\frac{1}{2(k-1)}} = \frac{1}{-\frac{1}{(k-1)}} = -(k-1)$$

Therefore,

$$\begin{aligned} \frac{d^2\Phi}{d\sigma^2} + 2\frac{k}{2}\frac{d\Phi}{d\sigma} + 0 + \{G^{11}\frac{\partial U(\Phi, \sigma)}{\partial \Phi} - [-(k-1)]G^{22}\frac{\partial U(\Phi, \sigma)}{\partial \sigma}\} &= 0 \\ \frac{d^2\Phi}{d\sigma^2} + k\frac{d\Phi}{d\sigma} + [G^{11}\frac{\partial U(\Phi, \sigma)}{\partial \Phi} + (k-1)G^{22}\frac{\partial U(\Phi, \sigma)}{\partial \sigma}] &= 0 \\ \frac{d^2\Phi}{d\sigma^2} + k\frac{d\Phi}{d\sigma} + \left[(e^{-\Phi-\sigma})\frac{\partial U(\Phi, \sigma)}{\partial \Phi} + (k-1)\left(\frac{1}{k-1}e^{-\Phi-\sigma}\right)\frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] &= 0 \\ \frac{d^2\Phi}{d\sigma^2} + k\frac{d\Phi}{d\sigma} + \left[(e^{-\Phi-\sigma})\frac{\partial U(\Phi, \sigma)}{\partial \Phi} + (e^{-\Phi-\sigma})\frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] &= 0 \\ \frac{d^2\Phi}{d\sigma^2} + k\frac{d\Phi}{d\sigma} + (e^{-\Phi-\sigma})\left[\frac{\partial U(\Phi, \sigma)}{\partial \Phi} + \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] &= 0 \end{aligned}$$

Potential function:

For case I: the equation of anharmonic system in a periodic field

$$U(\Phi, \sigma) = e^{\Phi+\sigma} \left[\frac{1}{2}\alpha\Phi^3 - \frac{3}{4}\alpha\Phi^2 - \frac{1}{4}(2\beta - 3\alpha)\Phi + \frac{1}{8}(2\beta - 3\alpha) - \frac{2b}{\omega^2+4}\cos(\omega\sigma) - \frac{\omega b}{\omega^2+4}\sin(\omega\sigma) \right]$$

$$\text{Let: } U(\Phi, \sigma) = e^{\Phi+\sigma} u(\Phi, \sigma)$$

$$\begin{aligned}
\frac{\partial U(\Phi, \sigma)}{\partial \Phi} &= e^{\Phi+\sigma} \left[u(\Phi, \sigma) + \frac{\partial u(\Phi, \sigma)}{\partial \Phi} \right] \\
&= e^{\Phi+\sigma} \left\{ \frac{1}{2} \alpha \Phi^3 - \frac{3}{4} \alpha \Phi^2 - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \Phi + \frac{1}{4} \left(\beta - \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right. \\
&\quad \left. + \frac{\partial \left[\frac{1}{2} \alpha \Phi^3 - \frac{3}{4} \alpha \Phi^2 - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \Phi + \frac{1}{4} \left(\beta - \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right]}{\partial \Phi} \right\} \\
&= e^{\Phi+\sigma} \left\{ \frac{1}{2} \alpha \Phi^3 - \frac{3}{4} \alpha \Phi^2 - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \Phi + \frac{1}{4} \left(\beta - \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right. \\
&\quad \left. + \left[\frac{3}{2} \alpha \Phi^2 - \frac{3}{2} \alpha \Phi - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \right] \right\} \\
&= e^{\Phi+\sigma} \left\{ \frac{1}{2} \alpha \Phi^3 - \frac{3}{4} \alpha \Phi^2 + \frac{3}{2} \alpha \Phi^2 - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \Phi - \frac{3}{2} \alpha \Phi + \frac{1}{4} \left(\beta - \frac{3}{2} \alpha \right) - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) \right. \\
&\quad \left. - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right\} = e^{\Phi+\sigma} \left\{ \frac{1}{2} \alpha \Phi^3 + \frac{3}{4} \alpha \Phi^2 - \frac{1}{2} \left(\beta + \frac{3}{2} \alpha \right) \Phi - \frac{1}{4} \left(\beta + \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right\}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial U(\Phi, \sigma)}{\partial \sigma} &= e^{\Phi+\sigma} \left[u(\Phi, \sigma) + \frac{\partial u(\Phi, \sigma)}{\partial \sigma} \right] \\
&= e^{\Phi+\sigma} \left\{ \frac{1}{2} \alpha \Phi^3 - \frac{3}{4} \alpha \Phi^2 - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \Phi + \frac{1}{4} \left(\beta - \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right. \\
&\quad \left. + \frac{\partial \left[\frac{1}{2} \alpha \Phi^3 - \frac{3}{4} \alpha \Phi^2 - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \Phi + \frac{1}{4} \left(\beta - \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right]}{\partial \sigma} \right\} \\
&= e^{\Phi+\sigma} \left\{ \frac{1}{2} \alpha \Phi^3 - \frac{3}{4} \alpha \Phi^2 - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \Phi + \frac{1}{4} \left(\beta - \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right. \\
&\quad \left. + \frac{\left[\partial \left[-\frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right] \right]}{\partial \sigma} \right\} \\
&= e^{\Phi+\sigma} \left\{ \frac{1}{2} \alpha \Phi^3 - \frac{3}{4} \alpha \Phi^2 - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \Phi + \frac{1}{4} \left(\beta - \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right. \\
&\quad \left. - \frac{\left[\partial \left[\frac{2b}{\omega^2 + 4} \cos(\omega\sigma) \right] + \partial \left[\frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right] \right]}{\partial \sigma} \right\} \\
&= e^{\Phi+\sigma} \left\{ \frac{1}{2} \alpha \Phi^3 - \frac{3}{4} \alpha \Phi^2 - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \Phi + \frac{1}{4} \left(\beta - \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) \right. \\
&\quad \left. - \left[-\frac{2b\omega}{\omega^2 + 4} \sin(\omega\sigma) + \frac{b\omega^2}{\omega^2 + 4} \cos(\omega\sigma) \right] \right\} \\
&= e^{\Phi+\sigma} \left\{ \frac{1}{2} \alpha \Phi^3 - \frac{3}{4} \alpha \Phi^2 - \frac{1}{2} \left(\beta - \frac{3}{2} \alpha \right) \Phi + \frac{1}{4} \left(\beta - \frac{3}{2} \alpha \right) - \frac{2b}{\omega^2 + 4} \cos(\omega\sigma) - \frac{\omega b}{\omega^2 + 4} \sin(\omega\sigma) + \frac{2b\omega}{\omega^2 + 4} \sin(\omega\sigma) \right\}
\end{aligned}$$

$$-\frac{b\omega^2}{\omega^2+4}\cos(\omega\sigma)\Big\}=e^{\Phi+\sigma}\left\{\frac{1}{2}\alpha\Phi^3-\frac{3}{4}\alpha\Phi^2-\frac{1}{2}\left(\beta-\frac{3}{2}\alpha\right)\Phi+\frac{1}{4}\left(\beta-\frac{3}{2}\alpha\right)-b\left(\frac{2+\omega^2}{\omega^2+4}\right)\cos(\omega\sigma)+\frac{b\omega}{\omega^2+4}\sin(\omega\sigma)\right\}$$

Therefore,

$$\begin{aligned}
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + (e^{-\Phi-\sigma}) \left[\frac{\partial U(\Phi, \sigma)}{\partial \Phi} + \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] = 0 \\
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + (e^{-\Phi-\sigma}) \left[e^{\Phi+\sigma} \left\{ \frac{1}{2}\alpha\Phi^3 + \frac{3}{4}\alpha\Phi^2 - \frac{1}{2}\left(\beta + \frac{3}{2}\alpha\right)\Phi - \frac{1}{4}\left(\beta - \frac{3}{2}\alpha\right) - \frac{2b}{\omega^2+4}\cos(\omega\sigma) - \frac{\omega b}{\omega^2+4}\sin(\omega\sigma) \right\} \right. \\
& \quad \left. + [e^{\Phi+\sigma} \left\{ \frac{1}{2}\alpha\Phi^3 - \frac{3}{4}\alpha\Phi^2 - \frac{1}{2}\left(\beta - \frac{3}{2}\alpha\right)\Phi + \frac{1}{4}\left(\beta - \frac{3}{2}\alpha\right) - b\left(\frac{2+\omega^2}{\omega^2+4}\right)\cos(\omega\sigma) + \frac{b\omega}{\omega^2+4}\sin(\omega\sigma) \right\}] \right] = 0 \\
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\frac{1}{2}\alpha\Phi^3 + \frac{3}{4}\alpha\Phi^2 - \frac{1}{2}\left(\beta + \frac{3}{2}\alpha\right)\Phi - \frac{1}{4}\left(\beta - \frac{3}{2}\alpha\right) - \frac{2b}{\omega^2+4}\cos(\omega\sigma) - \frac{\omega b}{\omega^2+4}\sin(\omega\sigma) \right] + \left[\frac{1}{2}\alpha\Phi^3 - \frac{3}{4}\alpha\Phi^2 - \frac{1}{2}\left(\beta - \frac{3}{2}\alpha\right)\Phi + \frac{1}{4}\left(\beta - \frac{3}{2}\alpha\right) + \frac{b\omega}{\omega^2+4}\sin(\omega\sigma) \right] = 0 \\
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left(\beta - \frac{3}{2}\alpha \right) - b\left(\frac{2+\omega^2}{\omega^2+4}\right)\cos(\omega\sigma) + \frac{b\omega}{\omega^2+4}\sin(\omega\sigma) = 0 \\
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \frac{1}{2}\alpha\Phi^3 + \frac{1}{4}\alpha\Phi^2 - \frac{3}{2}\left(\beta + \frac{3}{2}\alpha\right)\Phi - \frac{1}{4}\left(\beta - \frac{3}{2}\alpha\right)\Phi - \frac{1}{4}\left(\beta - \frac{3}{2}\alpha\right) + \frac{1}{4}\left(\beta - \frac{3}{2}\alpha\right) - \frac{2b}{\omega^2+4}\cos(\omega\sigma) \\
& \quad - b\left(\frac{2+\omega^2}{\omega^2+4}\right)\cos(\omega\sigma) - \frac{\omega b}{\omega^2+4}\sin(\omega\sigma) + \frac{b\omega}{\omega^2+4}\sin(\omega\sigma) = 0 \\
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \alpha\Phi^3 - \frac{1}{2}(2\beta)\Phi - \frac{4b+b\omega^2}{\omega^2+4}\cos(\omega\sigma) = 0 \\
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \alpha\Phi^3 - \beta\Phi - b\cos(\omega\sigma) = 0 \\
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \alpha\Phi^3 - \beta\Phi = b\cos(\omega\sigma)
\end{aligned}$$

For case II: the nonlinear equation of a parametrically excited damped pendulum

$$\begin{aligned} u(\Phi, \sigma) = & \frac{2}{5}\alpha \sin(\Phi) - \frac{1}{5}\alpha \cos(\Phi) \\ & + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) \\ & - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi)] \end{aligned}$$

$$\begin{aligned}
\frac{\partial U(\Phi, \sigma)}{\partial \Phi} &= e^{\Phi+\sigma} \left[u(\Phi, \sigma) + \frac{\partial u(\Phi, \sigma)}{\partial \Phi} \right] \\
&= e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right. \\
&\quad \left. + -\frac{\beta}{25+6\omega^2+\omega^4} [\omega(\omega^2+3) \sin(\omega\sigma) \sin(\Phi) - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2+5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2-5) \cos(\omega\sigma) \cos(\Phi)] \right] \\
&+ \frac{\partial \left(\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) + \frac{\beta}{25+6\omega^2+\omega^4} [\omega(\omega^2+3) \sin(\omega\sigma) \sin(\Phi) - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2+5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2-5) \cos(\omega\sigma) \cos(\Phi)] \right)}{\partial \Phi} \\
&= \\
&= e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right. \\
&\quad \left. + \frac{\beta}{25+6\omega^2+\omega^4} [\omega(\omega^2+3) \sin(\omega\sigma) \sin(\Phi) - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2+5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2-5) \cos(\omega\sigma) \cos(\Phi)] \right. \\
&\quad \left. + \frac{\partial \left(\frac{2}{5} \alpha \sin(\Phi) \right)}{\partial \Phi} + \frac{\partial \left(-\frac{1}{5} \alpha \cos(\Phi) \right)}{\partial \Phi} \right. \\
&\quad \left. + \frac{\beta}{25+6\omega^2+\omega^4} \frac{\partial \left([\omega(\omega^2+3) \sin(\omega\sigma) \sin(\Phi) - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2+5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2-5) \cos(\omega\sigma) \cos(\Phi)] \right)}{\partial \Phi} \right] \\
&= e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right. \\
&\quad \left. + \frac{\beta}{25+6\omega^2+\omega^4} [\omega(\omega^2+3) \sin(\omega\sigma) \sin(\Phi) - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2+5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2-5) \cos(\omega\sigma) \cos(\Phi)] \right]
\end{aligned}$$

$$\begin{aligned}
& + \frac{\partial \left(\frac{2}{5} \alpha \sin(\Phi) \right)}{\partial \Phi} + \frac{\partial \left(-\frac{1}{5} \alpha \cos(\Phi) \right)}{\partial \Phi} + \frac{\beta}{25 + 6\omega^2 + \omega^4} \left[\frac{\partial ([\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi)])}{\partial \Phi} - \frac{\partial ([4\omega \sin(\omega\sigma) \cos(\Phi)])}{\partial \Phi} \right. \\
& \left. + \frac{\partial (2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi))}{\partial \Phi} + \frac{\partial ((\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi))}{\partial \Phi} \right] \\
& = e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right. \\
& \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) \right. \\
& \left. - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi)] + \frac{2}{5} \alpha \cos(\Phi) + \frac{1}{5} \alpha \sin(\Phi) \right. \\
& \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \cos(\Phi) + 4\omega \sin(\omega\sigma) \sin(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi) \right. \\
& \left. - (\omega^2 - 5) \cos(\omega\sigma) \sin(\Phi)] \right] \\
& = e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) + \frac{2}{5} \alpha \cos(\Phi) + \frac{1}{5} \alpha \sin(\Phi) \right. \\
& \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) \right. \\
& \left. - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi) + \omega(\omega^2 + 3) \sin(\omega\sigma) \cos(\Phi) \right. \\
& \left. + 4\omega \sin(\omega\sigma) \sin(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi) - (\omega^2 - 5) \cos(\omega\sigma) \sin(\Phi)] \right] \\
& = e^{\Phi+\sigma} \left[\frac{3}{5} \alpha \sin(\Phi) + \frac{1}{5} \alpha \cos(\Phi) \right. \\
& \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) + 4\omega \sin(\omega\sigma) \sin(\Phi) \right. \\
& \left. - 4\omega \sin(\omega\sigma) \cos(\Phi) + \omega(\omega^2 + 3) \sin(\omega\sigma) \cos(\Phi) \right]
\end{aligned}$$

$$\begin{aligned}
& + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) - (\omega^2 - 5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi) \Big] \\
& = e^{\Phi+\sigma} \left[\frac{3}{5} \alpha \sin(\Phi) + \frac{1}{5} \alpha \cos(\Phi) + \frac{\beta}{25 + 6\omega^2 + \omega^4} [(\omega^3 + 7\omega) \sin(\omega\sigma) \sin(\Phi) + (\omega^3 - \omega) \sin(\omega\sigma) \cos(\Phi) \right. \\
& \left. + (\omega^2 + 15) \cos(\omega\sigma) \sin(\Phi) + (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi)] \right]
\end{aligned}$$

$$\begin{aligned}
\frac{\partial U(\Phi, \sigma)}{\partial \sigma} &= e^{\Phi+\sigma} \left[u(\Phi, \sigma) + \frac{\partial u(\Phi, \sigma)}{\partial \sigma} \right] = \\
&= e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right. \\
&\quad \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi)] \right. \\
&\quad \left. + \frac{\partial \left(\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi)] \right)}{\partial \sigma} \right. \\
&\quad \left. = e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right. \right. \\
&\quad \left. \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi)] \right. \right. \\
&\quad \left. \left. + \frac{\partial \left(\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi)] \right)}{\partial \sigma} \right. \right. \\
&\quad \left. \left. = e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right. \right. \right. \\
&\quad \left. \left. \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi)] \right. \right. \right]
\end{aligned}$$

$$\begin{aligned}
& + \frac{\partial \left(\frac{2}{5} \alpha \sin(\Phi) \right)}{\partial \sigma} + \frac{\partial \left(-\frac{1}{5} \alpha \cos(\Phi) \right)}{\partial \sigma} + \frac{\beta}{25 + 6\omega^2 + \omega^4} \left[\frac{\partial ([\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi))]}{\partial \sigma} - \frac{\partial ([4\omega \sin(\omega\sigma) \cos(\Phi))]}{\partial \sigma} \right. \\
& + \frac{\partial (2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi))}{\partial \sigma} + \left. \frac{\partial ((\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi))}{\partial \sigma} \right] \\
& = e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) + \frac{1}{5} \alpha \cos(\Phi) \right. \\
& \quad \left. - \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \sin(\omega\sigma) \sin(\Phi) \right. \\
& \quad \left. - 4\omega \sin(\omega\sigma) \cos(\Phi) + 2(\omega^2 + 5) \cos(\omega\sigma) \sin(\Phi) + (\omega^2 - 5) \cos(\omega\sigma) \cos(\Phi)] \right. \\
& \quad \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [\omega(\omega^2 + 3) \omega \cos(\omega\sigma) \sin(\Phi) - 4\omega^2 \cos(\omega\sigma) \cos(\Phi) - 2(\omega^2 + 5) \omega \sin(\omega\sigma) \square \sin(\Phi) \right. \\
& \quad \left. - (\omega^2 - 5) \omega \sin(\omega\sigma) \cos(\Phi)] \right] \\
& = e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right. \\
& \quad \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [-2(\omega^2 + 5) + (\omega^2 + 3) \omega \sin(\omega\sigma) \sin(\Phi) - (\omega^2 - 1) \omega \sin(\omega\sigma) \cos(\Phi) \right. \\
& \quad \left. + [\omega^2(\omega^2 + 3) + 2(\omega^2 + 5)] \cos(\omega\sigma) \sin(\Phi) - (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi)] \right] \\
& = e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right. \\
& \quad \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [-(\omega^2 + 7) \omega \sin(\omega\sigma) \sin(\Phi) - (\omega^2 - 1) \omega \sin(\omega\sigma) \cos(\Phi) + [\omega^4 + 5\omega^2 + 10] \cos(\omega\sigma) \sin(\Phi) \right. \\
& \quad \left. - (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi)] \right]
\end{aligned}$$

Thus,

$$\begin{aligned}
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + (e^{-\Phi-\sigma}) \left[\frac{\partial U(\Phi, \sigma)}{\partial \Phi} + \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] = 0 \\
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + (e^{-\Phi-\sigma}) \left[e^{\Phi+\sigma} \left[\frac{3}{5} \alpha \sin(\Phi) + \frac{1}{5} \alpha \cos(\Phi) \right] + \frac{\beta}{25 + 6\omega^2 + \omega^4} [(\omega^3 + 7\omega) \sin(\omega\sigma) \sin(\Phi) + (\omega^3 - \omega) \sin(\omega\sigma) \cos(\Phi) \right. \right. \\
& \quad \left. \left. + (\omega^2 + 15) \cos(\omega\sigma) \sin(\Phi) + (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi)] \right] \right. \\
& \quad \left. + e^{\Phi+\sigma} \left[\frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right] \right. \\
& \quad \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [-(\omega^2 + 7) \omega \sin(\omega\sigma) \sin(\Phi) - (\omega^2 - 1) \omega \sin(\omega\sigma) \cos(\Phi) + [\omega^4 + 5\omega^2 + 10] \cos(\omega\sigma) \sin(\Phi) \right. \\
& \quad \left. - (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi)] \right] = 0 \\
& \frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\frac{3}{5} \alpha \sin(\Phi) + \frac{1}{5} \alpha \cos(\Phi) \right. \\
& \quad \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [(\omega^3 + 7\omega) \sin(\omega\sigma) \sin(\Phi) + (\omega^3 - \omega) \sin(\omega\sigma) \cos(\Phi) + (\omega^2 + 15) \cos(\omega\sigma) \sin(\Phi) \right. \\
& \quad \left. + (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi)] + \frac{2}{5} \alpha \sin(\Phi) - \frac{1}{5} \alpha \cos(\Phi) \right. \\
& \quad \left. + \frac{\beta}{25 + 6\omega^2 + \omega^4} [-(\omega^2 + 7) \omega \sin(\omega\sigma) \sin(\Phi) - (\omega^2 - 1) \omega \sin(\omega\sigma) \cos(\Phi) + [\omega^4 + 5\omega^2 + 10] \cos(\omega\sigma) \sin(\Phi) \right. \\
& \quad \left. - (3\omega^2 + 5) \cos(\omega\sigma) \cos(\Phi)] \right] = 0
\end{aligned}$$

$$\begin{aligned}
\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\frac{3}{5}\alpha \sin(\Phi) + \frac{1}{5}\alpha \cos(\Phi) + \frac{2}{5}\alpha \sin(\Phi) - \frac{1}{5}\alpha \cos(\Phi) + \frac{\beta}{25+6\omega^2+\omega^4} [-(\omega^2+7)\omega \sin(\omega\sigma) \sin(\Phi) \right. \\
\left. - (\omega^2-1)\omega \sin(\omega\sigma) \cos(\Phi) + [\omega^4+5\omega^2+10]\cos(\omega\sigma) \sin(\Phi) - (3\omega^2+5)\cos(\omega\sigma) \cos(\Phi) + (\omega^3+7\omega)\sin(\omega\sigma) \sin(\Phi) \right. \\
\left. + (\omega^3-\omega)\sin(\omega\sigma) \cos(\Phi) + (\omega^2+15)\cos(\omega\sigma) \sin(\Phi) + (3\omega^2+5)\cos(\omega\sigma) \cos(\Phi)] \right] = 0
\end{aligned}$$

$$\begin{aligned}
\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25+6\omega^2+\omega^4} [-(\omega^2+7)\omega \sin(\omega\sigma) \sin(\Phi) - (\omega^2-1)\omega \sin(\omega\sigma) \cos(\Phi) + [\omega^4+5\omega^2+10]c \square s(\omega\sigma) \sin(\Phi) \right. \\
\left. - (3\omega^2+5)\cos(\omega\sigma) \cos(\Phi) + (\omega^3+7\omega)\sin(\omega\sigma) \sin(\Phi) + (\omega^3-\omega)\sin(\omega\sigma) \cos(\Phi) + (\omega^2+15)\cos(\omega\sigma) \sin(\Phi) \right. \\
\left. + (3\omega^2+5)\cos(\omega\sigma) \cos(\Phi) + \frac{2}{5}\alpha \sin(\Phi) - \frac{1}{5}\alpha \cos(\Phi)] \right] = 0
\end{aligned}$$

$$\begin{aligned}
\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25+6\omega^2+\omega^4} [-(\omega^2+7)\omega \sin(\omega\sigma) \sin(\Phi) + (\omega^3+7\omega)\sin(\omega\sigma) \sin(\Phi) - (\omega^2-1)\omega \sin(\omega\sigma) \cos(\Phi) \right. \\
\left. + (\omega^3-\omega)\sin(\omega\sigma) \cos(\Phi) + [\omega^4+5\omega^2+10]\cos(\omega\sigma) \sin(\Phi) + (\omega^2+15)\cos(\omega\sigma) \sin(\Phi) - (3\omega^2+5)\cos(\omega\sigma) \cos(\Phi) \right. \\
\left. + (3\omega^2+5)\cos(\omega\sigma) \cos(\Phi)] \right] = 0
\end{aligned}$$

$$\begin{aligned}
\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25+6\omega^2+\omega^4} [-(\omega^2+7)\omega \sin(\omega\sigma) \sin(\Phi) + (\omega^3+7\omega)\sin(\omega\sigma) \sin(\Phi) - (\omega^2-1)\omega \sin(\omega\sigma) \cos(\Phi) \right. \\
\left. + (\omega^3-\omega)\sin(\omega\sigma) \cos(\Phi) + [\omega^4+5\omega^2+10]\cos(\omega\sigma) \sin(\Phi) + (\omega^2+15)\cos(\omega\sigma) \sin(\Phi) - (3\omega^2+5)\cos(\omega\sigma) \cos(\Phi) \right. \\
\left. + (3\omega^2+5)\cos(\omega\sigma) \cos(\Phi)] \right] = 0
\end{aligned}$$

$$\begin{aligned}
\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25+6\omega^2+\omega^4} [-(\omega^2+7)\omega \sin(\omega\sigma) \sin(\Phi) + (\omega^3+7\omega)\sin(\omega\sigma) \sin(\Phi) - (\omega^2-1)\omega \sin(\omega\sigma) \cos(\Phi) + (\omega^3- \\
\omega)\sin(\omega\sigma) \cos(\Phi) + [\omega^4+5\omega^2+10]\cos(\omega\sigma) \sin(\Phi) + (\omega^2+15)\cos(\omega\sigma) \sin(\Phi) - (3\omega^2+5)\cos(\omega\sigma) \cos(\Phi) + \\
(3\omega^2+5)\cos(\omega\sigma) \cos(\Phi)] \right] = 0
\end{aligned}$$

$$\begin{aligned}
\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25+6\omega^2+\omega^4} [(\omega^4+5\omega^2+10+\omega^2+15)\cos \square (\omega\sigma) \sin(\Phi)] \right] = 0
\end{aligned}$$

$$\begin{aligned}
\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \left[\alpha \sin(\Phi) + \frac{\beta}{25+6\omega^2+\omega^4} [(\omega^4+6\omega^2+25)\cos(\omega\sigma) \sin(\Phi)] \right] = 0
\end{aligned}$$

$$\begin{aligned}
\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + \alpha \sin(\Phi) + \beta \cos(\omega\sigma) \sin(\Phi) = 0
\end{aligned}$$

$$\begin{aligned}
\frac{d^2\Phi}{d\sigma^2} + k \frac{d\Phi}{d\sigma} + [\alpha + \beta \cos(\omega\sigma)] \sin(\Phi) = 0
\end{aligned}$$

For case III: the equation of anharmonic system in a periodic field

$$\begin{aligned}
 \frac{d^2\Phi}{d\sigma^2} + (e^{-\Phi-\sigma}) \left[\frac{\partial U(\Phi, \sigma)}{\partial \Phi} + \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] &= 0 \\
 U(\Phi, \sigma) &= e^{\Phi+\sigma} \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 - \frac{m}{4\hbar^2} K\sigma^2 - \frac{m}{2\hbar^2} K\Phi\sigma + \frac{2m}{\hbar^2} (E - K)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{\hbar^2} \left(\frac{3}{4}K - E\right) \right] \\
 U(\Phi, \sigma) &= e^{\Phi+\sigma} \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} \sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} K\sigma + \frac{m}{2\hbar^2} \left(\frac{3}{4}K - E\right) \right]
 \end{aligned}$$

$$\begin{aligned}
\frac{\partial U(\Phi, \sigma)}{\partial \Phi} &= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} (\frac{3}{4}K - E) \right. \\
&\quad \left. + \frac{\partial \left[-\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} (\frac{3}{4}K - E) \right]}{\partial \Phi} \right\} \\
&= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{2m}{\hbar^2} (E - K) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{\hbar^2} (\frac{3}{4}K - E) + \frac{\partial \left[-\frac{m}{2\hbar^2} K \Phi \sigma^2 \right]}{\partial \Phi} + \frac{\partial \left[\frac{m}{4\hbar^2} K \sigma^2 \right]}{\partial \Phi} \right\} \\
&\quad + \frac{\partial \left[\frac{m}{2\hbar^2} K \Phi \sigma \right]}{\partial \Phi} + \frac{\partial \left[+\frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi \right]}{\partial \Phi} + \frac{\partial \left[-\frac{m}{2\hbar^2} K \sigma \right]}{\partial \Phi} + \frac{\partial \left[+\frac{m}{2\hbar^2} (\frac{3}{4}K - E) \right]}{\partial \Phi} \Big\} = \\
&= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} (\frac{3}{4}K - E) - \frac{m}{2\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \sigma \right. \\
&\quad \left. + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \right\} \\
&= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 - \frac{m}{2\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} (\frac{3}{4}K - E) \right. \\
&\quad \left. + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \right\} = e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 - \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi + \frac{m}{2\hbar^2} (\frac{1}{4}K + E) \right\}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial U(\Phi, \sigma)}{\partial \sigma} &= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} (\frac{3}{4}K - E) \right. \\
&\quad \left. + \frac{\partial \left[-\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} (\frac{3}{4}K - E) \right]}{\partial \sigma} \right\} \\
&= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} (\frac{3}{4}K - E) + \frac{\partial \left[-\frac{m}{2\hbar^2} K \Phi \sigma^2 \right]}{\partial \sigma} \right. \\
&\quad \left. + \frac{\partial \left[\frac{m}{4\hbar^2} K \sigma^2 \right]}{\partial \sigma} + \frac{\partial \left[\frac{m}{2\hbar^2} K \Phi \square \right]}{\partial \sigma} + \frac{\partial \left[\frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi \right]}{\partial \sigma} + \frac{\partial \left[-\frac{m}{2\hbar^2} K \sigma \right]}{\partial \sigma} + \frac{\partial \left[\frac{m}{2\hbar^2} (\frac{3}{4}K - E) \right]}{\partial \sigma} \right\} \\
&= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} (\frac{3}{4}K - E) - \frac{m}{2\hbar^2} K \Phi \frac{\partial[\sigma^2]}{\partial \sigma} + \frac{m}{4\hbar^2} K \frac{\partial[\sigma^2]}{\partial \sigma} \right. \\
&\quad \left. + \frac{m}{2\hbar^2} K \Phi \frac{\partial[\square]}{\partial \sigma} + 0 - \frac{m}{2\hbar^2} K \frac{\partial[\sigma]}{\partial \sigma} + 0 \right\} \\
&= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} (\frac{3}{4}K - E) - \frac{m}{\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} K \Phi \right. \\
&\quad \left. - \frac{m}{2\hbar^2} K \right\} \\
&= e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E - \frac{1}{2}K) \Phi + \frac{m}{2\hbar^2} K \sigma - \frac{m}{2\hbar^2} K \Phi - \frac{m}{2\hbar^2} K \sigma + \frac{m}{2\hbar^2} (\frac{3}{4}K - E) \right. \\
&\quad \left. - \frac{m}{2\hbar^2} K \right\} = e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K \Phi \sigma^2 + \frac{m}{4\hbar^2} K \sigma^2 + \frac{m}{2\hbar^2} K \Phi \sigma + \frac{m}{2\hbar^2} (2E + \frac{1}{2}K) \Phi - \frac{m}{2\hbar^2} (\frac{1}{4}K + E) \right\} \\
&\quad \frac{d^2 \Phi}{d\sigma^2} + (e^{-\Phi-\sigma}) \left[\frac{\partial U(\Phi, \sigma)}{\partial \Phi} + \frac{\partial U(\Phi, \sigma)}{\partial \sigma} \right] = 0
\end{aligned}$$

$$\begin{aligned} \frac{d^2\Phi}{d\sigma^2} + (e^{-\Phi-\sigma}) \left[e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K\Phi\sigma^2 - \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi + \frac{m}{2\hbar^2} \left(\frac{1}{4}K + E\right) \right\} \right. \\ \left. + e^{\Phi+\sigma} \left\{ -\frac{m}{2\hbar^2} K\Phi\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 - \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E + \frac{1}{2}K\right)\Phi - \frac{m}{2\hbar^2} \left(\frac{1}{4}K + E\right) \right\} \right] = 0 \end{aligned}$$

$$\begin{aligned} \frac{d^2\Phi}{d\sigma^2} + \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 - \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi + \frac{m}{2\hbar^2} \left(\frac{1}{4}K + E\right) \right. \\ \left. - \frac{m}{2\hbar^2} \left(\frac{1}{4}K + E\right) \right] = 0 \end{aligned}$$

$$\begin{aligned} \frac{d^2\Phi}{d\sigma^2} + \left[-\frac{m}{2\hbar^2} K\Phi\sigma^2 - \frac{m}{2\hbar^2} K\Phi\sigma^2 - \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{4\hbar^2} K\sigma^2 + \frac{m}{2\hbar^2} K\Phi\sigma - \frac{m}{2\hbar^2} K\Phi\sigma + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K\right)\Phi + \frac{m}{2\hbar^2} \left(\frac{1}{4}K + E\right) \right. \\ \left. - \frac{m}{2\hbar^2} \left(\frac{1}{4}K + E\right) \right] = 0 \end{aligned}$$

$$\frac{d^2\Phi}{d\sigma^2} + \left[-\frac{m}{\hbar^2} K\Phi\sigma^2 + \frac{m}{2\hbar^2} \left(2E - \frac{1}{2}K + 2E + \frac{1}{2}K\right)\Phi \right] = 0$$

$$\frac{d^2\Phi}{d\sigma^2} + \left[-\frac{m}{\hbar^2} K\Phi\sigma^2 + \frac{2m}{\hbar^2} E\Phi \right] = 0$$

$$\frac{d^2\Phi}{d\sigma^2} + \left[-\frac{m}{\hbar^2} K\Phi\sigma^2 + \frac{2m}{\hbar^2} E\Phi \right] = 0$$

$$\frac{d^2\Phi}{d\sigma^2} + \frac{2m}{\hbar^2} \left[-\frac{1}{2} K\Phi\sigma^2 + E\Phi \right] = 0$$

$$\frac{d^2\Phi}{d\sigma^2} + \frac{2m}{\hbar^2} \left[+E - \frac{1}{2} K\sigma^2 \right] \Phi = 0$$

$$\frac{d^2\Phi}{d\sigma^2}=-\frac{2m}{\hbar^2}\left(E-\frac{1}{2}K\sigma^2\right)\phi(\sigma)\,,$$